

File Name	36787_41534_25kg_MPL_VO2.pdf
Dimensions	390 x 140 x 1010mm
Scale	A3 = 26%
Date	01/10/2007



1010

178mm

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY



ACTIVE CONSTITUENT: 100g/kg MONENSIN (as monensin sodium)

- Feedlot Cattle:** For improved feed efficiency. As an aid in the control of bloat.
- Pasture Cattle:** For improved weight gain and feed efficiency.
- Dairy cows:** For increased milk production under many conditions. As an aid in reducing the severity of non-clinical ketosis in lactating dairy cows. As an aid in the control of bloat.
- Cattle:** For improved weight gain and feed efficiency in heifers. For improved reproductive performance in heifers. As an aid in the prevention of coccidiosis caused by *Eimeria zuernii* and *Eimeria bovis*.
- Goats:** As an aid in the prevention of coccidiosis.
- Chickens:** As an aid in the prevention of coccidiosis caused by *Eimeria acervulina*, *E. brunetti*, *E. maxima*, *E. mivati*, *E. necatrix* and *E. tenella*. (For use in broiler and layer replacement chicken feeds only).

Store below 30°C (Room Temperature) in a dry place.
For expiry date and batch number see side of bag.

APVMA Approval No: 36787/25kg/0107

ELANCO ANIMAL HEALTH
A DIVISION OF ELI LILLY AUSTRALIA PTY LIMITED
A.B.N. 39 000 233 992
WHARF ROAD, WEST RYDE, N.S.W. 2114
Tel: Toll Free 1800 226 324

25kg NET



* Elanco® Elancoban® and the diagonal colour bar are trademarks of Eli Lilly and Company

USE NO HOOKS



125mm

Expiry Date:
Batch No:
BG7033AALB



ELANCO® AF1304 Elancoban G Monensin Sodium Premix

DIRECTIONS FOR USE:
Restraints:
DO NOT USE in chickens producing eggs.
DO NOT USE in goats producing milk for human consumption.
Contraindications:
Dogs, horses and other equines must not be allowed access to feeds containing Elancoban G. Ingestion of Elancoban G may be fatal in these species.
Precautions:
The recommended levels of Elancoban G Premix must not be exceeded. Excessively high levels due to mixing errors can cause reduced feed intake, poor growth and diarrhoea, and death. Poultry consuming monensin sodium should not be treated with tiamulin. Severe growth depression or death may occur. Responses to Elancoban G Premix are unlikely if nutrient intake is significantly restricted. Responses to Elancoban G Premix in pasture cattle may not occur if pasture quality is poor or if weather conditions are very hot and dry.
Mixing Directions:
IMPORTANT - THIS PREMIX MUST BE THOROUGHLY MIXED WITH THE OTHER FEED COMPONENTS BEFORE USE AND CARE SHOULD BE TAKEN IN TRANSPORTING MIXING FEED TO PREVENT SETTLING.

	Complete Feed ¹		Supplements	
	Monensin ppm in feed ²	Elancoban G kg/tonne of feed	Monensin mg/head/day	Elancoban G grams/head/day
CATTLE				
Feed Efficiency	11 - 33	0.11 - 0.33	50 - 360	0.5 - 3.6
Feed Efficiency and Weight Gain (Pasture Cattle) ³			100 - 200 ⁴	1.0 - 2.0
Weight Gain, Reproductive Performance (Heifers)	11 - 33	0.11 - 0.33	200 - 360	2.0 - 3.6
Coccidiosis (for DAIRY COWS see section below)	11 - 33	0.11 - 0.33		
Bloat (for DAIRY COWS see section below)	25 - 33	0.25 - 0.33		
DAIRY COWS⁵				
Milk Production, Ketosis	6 - 18	0.06 - 0.18	200 - 450	2.0 - 4.5
Coccidiosis	11 - 18	0.11 - 0.18	200 - 450	2.0 - 4.5
Bloat				
GOATS				
Coccidiosis	5 - 20	0.05 - 0.20	7.5 - 40.0	0.075 - 0.40
CHICKENS (Broilers and Replacement Pullets)				
Coccidiosis	100 - 120	1.0 - 1.2		
	Monensin ppm in feed	Elancoban G kg/tonne of feed	Monensin mg/kg BW/day ⁶	Elancoban G mg/kg BW/day
CALVES (50 - 250 kg)	11 - 33	0.11 - 0.33	0.5 - 1.0	5.0 - 15.0
Coccidiosis				

¹ Calculated on a 90% dry matter basis. Rations containing silage or other wet feeds should be corrected to a 90% dry matter basis for calculation of the appropriate monensin dosage.
² ppm = parts per million which is equivalent to mg/kg.
³ Feeding monensin to lactating dairy cows may result in decreased milk fat percentage, but there is no effect on total fat yield.
⁴ kg BW = kilograms bodyweight.
⁵ For example, cattle supplement containing Elancoban G should be fed at 0.06kg/head, divided into morning and evening feeds.
⁶ The optimum dose is 20mg monensin/head/day.

GENERAL MIXING INSTRUCTIONS:
IMPORTANT: Thoroughly mixing Elancoban G into an intermediate premix (e.g. 25kg) and mixing the intermediate into the finished feed is recommended to ensure homogeneity.

1. COMPLETE FEEDS: The following are examples of the amount of Elancoban G to be incorporated into 1 tonne (1000 kg) of complete feed for various levels.

Concentration of monensin required in complete feed (ppm or mg/kg)	Amount of Elancoban G Premix (kg) to be added to 1 tonne of feed
5	0.05
6	0.06
11	0.11
18	0.18
20	0.20
33	0.33

2. SUPPLEMENTS: For supplements, adjust the quantity of Elancoban G according to the formula below. Do not feed a supplement containing greater than 300 mg monensin/kg supplement. The supplement feeding level may need to be adjusted to ensure animals receive the correct daily dose of monensin.

$$\text{Kilograms of Elancoban G to add to 1 tonne of feed} = \frac{\text{milligrams of monensin required per animal per day} \times 100}{\text{kilograms of supplement fed per animal per day}}$$

Feeding Directions:
Broilers - Feed continuously as the only ration.
Layer Replacements - Feed only to birds that will be placed into cages and feed until they are placed into cages.

WITHHOLDING PERIODS:
Cattle: MEAT: NIL
MILK: NIL
Goats: MEAT: NIL
MILK: DO NOT USE Elancoban G in goats producing milk for human consumption.
Chickens: NIL

SAFETY DIRECTIONS: Poisonous if swallowed. Will irritate the eyes, nose, throat and skin. Avoid contact with eyes and skin. Repeated exposure may cause allergic disorders. Do not inhale dust when opening the container and mixing into feed. Wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves, and half facemask respirator with dust cartridge or canister. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, respirator and if rubber wash with detergent and warm water, and contaminated clothing.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 13 1126. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

DISPOSAL OF BAGS: Shake and empty contents into medicated feed. Do not dispose of undiluted chemicals on site. Puncture or shred and bury empty bags in a local authority landfill. If not available bury the bag below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty bags and product should not be burnt.

Additional information is in the Material Safety Data Sheet.