

**POISON**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**  
**FOR ANIMAL TREATMENT ONLY**



Anticoccidial Premix



**ACTIVE CONSTITUENTS:** NARASIN 80g/kg  
NICARBAZIN 80g/kg

For the prevention of coccidiosis caused by *Eimeria acervulina*, *E. brunetti*, *E. maxima*, *E. mivati*, *E. necatrix*, and *E. tenella*.

FOR USE IN BROILER CHICKEN FEEDS ONLY.

**25 kg NET**

**DISPOSAL OF BAGS:** Shake and empty contents into medicated feed. Do not dispose of undiluted chemicals on site. Puncture or shred and bury empty bags in a local authority landfill. If not available bury the bag below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty bags and product should not be burnt.

Store below 30° (Room Temperature) in a dry place.  
For expiry date and batch number see side of bag

NRA Approval No. 36802/01

Member  
(logo) Avcare

ELANCO ANIMAL HEALTH  
A division of Eli Lilly Australia Pty Limited  
A.B.N. 39 000 233 992  
112 WHARF ROAD, WEST RYDE, N.S.W. 2114  
TEL: Toll Free 1800 226 324

## USE NO HOOKS

GENERIC PRODUCT NAME:	FILE NAME:	VERSION:	REVISION:	LABEL CODE	
Maxiban	36802_01c.doc	1	Created on 27/09/01 13:12	BG7093A ALB	Page 1 of 3

(Back of bag)

**DIRECTIONS FOR USE:**

**DO NOT USE in laying hens.**

**DO NOT USE in turkeys or game birds.**

**DO NOT USE in horses or other equines - may be fatal.**

**Poultry consuming narasin should not be treated with tiamulin (Dynamutilin®).**

**Severe growth depression or death may occur.**

**MIXING DIRECTIONS:**

**IMPORTANT - THIS PREMIX MUST BE THOROUGHLY MIXED IN FEED BEFORE USE.**

Thoroughly mix 0.5 to 0.625kg MAXIBAN Premix in one tonne of feed to provide 80 to 100 ppm of total active ingredient per tonne of complete feed.

**FEEDING DIRECTIONS:** Feed continuously as the only ration for at least 14 days.

**WITHHOLDING PERIOD: NIL**

**SAFETY DIRECTIONS: Poisonous if swallowed. Will damage eyes and will irritate the nose, throat and skin. Avoid contact with eyes and skin. Repeated exposure may cause allergic disorders. Do not inhale dust when opening the container and mixing into feed. Wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves, goggles and disposable dust mask. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.**

**FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 13 1126. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.**

Additional information is in the Material Safety Data Sheet.

GENERIC PRODUCT NAME:	FILE NAME:	VERSION:	REVISION:	LABEL CODE	
Maxiban	36802_01c.doc	1	Created on 27/09/01 13:12	BG7093A ALB	Page 2 of 3

Side Panel 1

**EJANCO** AF1375



Anticoccidial Premix

BG7093AALB

Side panel 2

(expiry date and batch number)

GENERIC PRODUCT NAME:	FILE NAME:	VERSION:	REVISION:	LABEL CODE	
Maxiban	36802_01c.doc	1	Created on 27/09/01 13:12	BG7093A ALB	Page 3 of 3