

READ SAFETY DIRECTIONS BEFORE OPENING OR USING FOR ANIMAL TREATMENT ONLY

ELANCO* AF0375

Surmax® 100

avilamycin

*



Premix

ACTIVE CONSTITUENT: 100g/kg AVILAMYCIN

For increased rate of weight gain and improved feed efficiency in broiler chickens.

As an aid in the prevention of necrotic enteritis due to *Clostridium perfringens* in broiler chickens.

DIRECTIONS FOR USE:

25 kg NET

**RESTRAINTS: DO NOT USE in feed for food animal species other than broiler chickens.
DO NOT USE in laying birds where eggs or egg products are to be used for
human consumption or processing.**

DOSAGE AND ADMINISTRATION:

Mixing and feeding directions:

Surmax 100 Premix must be thoroughly mixed in feeds before use and should be thoroughly mixed with a small quantity of feed ingredient (20 to 50kg) before it is incorporated into the bulk of the feed to be prepared.

BROILER CHICKENS:

For increased rate of weight gain and improved feed efficiency:

Thoroughly mix 25 to 150 grams of SURMAX 100 into 1000kg of feed to provide 2.5 to 15 grams avilamycin per tonne or 2.5 to 15 ppm. Feed continuously as the only ration.

As an aid in the prevention of necrotic enteritis:

Thoroughly mix 100 to 150 grams of SURMAX 100 into 1000 kg of feed to provide 10 to 15 grams avilamycin per tonne or 10 to 15 ppm. Feed continuously as the only ration during the period that birds are considered to be at risk of the disease.

Most outbreaks of necrotic enteritis occur in broilers from 2 to 6 weeks of age.

Appropriate additional measures to manage necrotic enteritis in broilers:

Clean and disinfect buildings between batches of birds, maintain dry litter, ensure proper ventilation, avoid overcrowding. Reduce immunosuppressive stress and disease. Use effective coccidiosis control. Evaluate bird nutrition, particularly protein and cereal sources and fibre content of feed. Comprehensive necrotic enteritis management should be planned in consultation with veterinary and nutritional poultry expertise.

WITHHOLDING PERIODS - MEAT: NIL

EGGS: DO NOT USE in laying birds where eggs or egg products are to be used for human consumption or processing.

An Export Slaughter Interval may apply to this product. Contact Elanco Animal Health before using this product in poultry destined for the export market.

SAFETY DIRECTIONS: May irritate the eyes and skin. Avoid contact with the eyes and skin. When opening the container and mixing into stock feed, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length PVC gloves, and a disposable dust mask. Wash hands after use. After each day's use, wash gloves and contaminated clothing.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 13 1126.

DISPOSAL OF BAGS: Shake and empty contents into medicated feed. Do not dispose of undiluted chemicals on site. Puncture or shred and bury empty bags in a local authority landfill. If not available bury the bag below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty bags and product should not be burned.

APVMA Approval No. 49376/25kg/0305

Manufactured in U.K. for:

ELANCO ANIMAL HEALTH

A DIVISION OF ELI LILLY AUSTRALIA PTY. LIMITED

A.B.N. 39 000 233 992

112 WHARF ROAD, WEST RYDE, N.S.W. 2114

TEL: (toll free) 1800 226 324

Store below 30°C (Room Temperature) in a dry place

MEMBER
Avicare



BG7201DEALL (I-FEB-05)
BGG120

Batch number:

Expiry date:

USE NO HOOKS

* Elanco®, Surmax® and the diagonal colour bar are trademarks of Eli Lilly and Company