


File Name	36791_37969_20kg_label_MPL_V01.pdf
Dimensions	390 x 140 x 910mm
Scale	A3 = 26% A4=18%
Date	24/07/2007



**CAUTION**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING  
FOR ANIMAL TREATMENT ONLY



ACTIVE CONSTITUENT: 50g/kg TYLOSIN as the phosphate

For use as an aid in stimulating growth and improving feed efficiency in pigs.

For the treatment and prevention of ileitis in pigs.

For reduction in the incidence of liver abscess in cattle.

As an aid in the control of necrotic enteritis caused by *Clostridium perfringens* in broiler and replacement chickens.

**20kg NET**

Store below 30°C (Room Temperature) in a dry place.  
For expiry date and batch number see side of bag.

APVMA Approval No: 36791/20/1106

Manufactured in Australia by:  
ELANCO ANIMAL HEALTH  
A DIVISION OF ELI LILLY AUSTRALIA PTY LIMITED  
A.B.N. 39 000 233 992  
112 WHARF ROAD, WEST RYDE, N.S.W. 2114  
Tel: Toll Free 1800 226 324



\* Elanco®, Tylan® and the diagonal colour bar are trademarks of Eli Lilly and Company

**USE NO HOOKS**



**BG780001AL**



ELANCO <sup>AF0050</sup> Tylan 50 Tylosin Phosphate Premix

**DIRECTIONS FOR USE:**  
Mixing directions:  
**IMPORTANT - TYLAN 50 PREMIX MUST BE THOROUGHLY MIXED IN FEED BEFORE USE.**

Recommended levels for use:

A. PIGS	Amount of Tylan 50 Premix mixed per tonne of feed	Tylosin Equivalent per tonne of feed
For stimulating growth and improving feed efficiency:		
a) Up to 20kg liveweight	2.0kg	100g
b) 20kg to 50kg liveweight	0.4kg	20g
c) 50kg to market	0.2kg	10g
For the prevention of ileitis:	0.8kg	40g

**B. CATTLE**

For reduction in the incidence of liver abscess:

In Complete Feeds: 11 ppm tylosin, i.e. 0.22kg Tylan 50 Premix per tonne of feed.  
Supplementary Feeds: Dependent on total dry matter intake of animal and supplementary feeding rate. Ensure that amount of Tylan fed does not exceed 11ppm of total dry matter intake (supplement plus all other feeds consumed per day). As an example, for dairy cows, addition according to the following table provides 150mg tylosin/head/day.

Supplementary Feeding Rate kg/head/day	Amount of Tylan 50 Premix per tonne of feed	Tylosin equivalent per tonne of feed
1	3.0 kg	150.0 g
2	1.5 kg	75.0 g
4	0.75 kg	37.5 g
6	0.5 kg	25.0 g
8	0.375 kg	18.75g

Make changes to diet gradually. Increase tylosin in feed up to desired dose over 5-7 days.

**C. POULTRY**

For control of necrotic enteritis.

Thoroughly mix Tylan 50 Premix in feed at a rate of 100 mg tylosin per kg (2.0kg TYLAN 50 Premix per tonne) of feed. Medicated feed should be provided as the sole ration to growing chickens for a period of 7 days when necrotic enteritis has been identified or can be predicted to occur.

**Additional measures to manage necrotic enteritis in broilers:**

Clean and disinfect buildings between batches of birds, maintain dry litter, ensure proper ventilation, avoid overcrowding. Reduce immunosuppressive stress and disease. Use effective coccidiosis control. Evaluate bird nutrition, particularly protein and cereal sources and fibre content of feed. Comprehensive necrotic enteritis management should be planned in consultation with veterinary and nutritional poultry expertise.

Any variation by the prescribing veterinarian to the approved dose, frequency, duration, route, disease or target species may require extending the approved withholding period.

**WITHOLDING PERIODS:**

MEAT (pigs, cattle, poultry): NIL

MILK: NIL

EGGS: NIL

**SAFETY DIRECTIONS:** Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. Repeated exposure may cause allergic disorders. Do not inhale dust. When opening the container and mixing into premises wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow length PVC gloves and disposable dust face mask covering mouth and nose. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

**FIRST AID:** If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 13 1126.

**DISPOSAL OF BAGS:** Shake and empty contents into medicated feed. Do not dispose of undiluted chemicals on site. Puncture or shred and bury empty bags in a local authority landfill. If not available bury the bag below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty bags and product should not be burned.

Additional information is in the Material Safety Data Sheet.