

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY



Premix

Active constituent: 20g/kg ractopamine hydrochloride (equivalent to 17.8 g/kg ractopamine)

For increased rate of weight gain, improved feed efficiency, increased carcass leanness and increased dressing percentage in finisher pigs.

10kg NET

DIRECTIONS FOR USE:

Restraint:

DO NOT ADMINISTER PAYLEAN TO SPECIES OTHER THAN PIGS.

Note to feed formulators:

Unintended inclusion of medication into animal feeds may result in residues that adversely impact trade access or the rules of professional competition. Use cleaning or flushing, or avoid sequencing feed for other species (in particular ruminant and horse feeds) immediately after manufacture of pig finisher feeds containing Paylean.

A copy of directions for use relevant to final feed are required to be provided to the producers with each consignment of Paylean treated feed. Contact Elanco Animal Health for further information.

**NOT FOR USE IN BREEDING PIGS.
FOR USE IN FINISHER PIG FEEDS ONLY.**

Dosage and Administration:

Recommended dosages of ractopamine hydrochloride:

- 5-20 ppm in final feed for increased rate of weight gain and improved feed efficiency
- 10-20 ppm in final feed for increased carcass leanness and increased carcass dressing percent

Product Name	File Name	Version	Revision Date	Page No.
Paylean 20	Paylean 20 54172 1205 for Infopest.doc	Created on 3/14/2006 4:27:00 PM	Last printed 3/14/2006 4:31:00 PM	Page 1 of 3

Mixing Directions:**PAYLEAN 20 PREMIX MUST BE THOROUGHLY MIXED INTO FEEDS BEFORE USE.**

All calculations are based on 90% dry matter.

Ractopamine Concentration in Final Feed g/tonne (ppm)	Grams of Paylean 20 Premix to Add per Tonne of Final Feed
5	250
10	500
15	750
20	1000 (1 kg)

IMPORTANT – To ensure adequate mixing, it is recommended that a pre-blend of Paylean 20 Premix be made before incorporation into the total amount of complete feed. Thoroughly mix the pre-blend into a nutritionally balanced pig ration to result in a final ration containing 5 to 20 ppm of ractopamine. Amino acid content of the ration may influence response to treatment. Dietary specifications should be determined in consultation with a recognised pig nutritionist in order to optimise Paylean benefits.

Feeding directions: Feed complete ration containing ractopamine continuously as the sole ration to finisher pigs for the last 28-42 days prior to slaughter.

Pigs fed Paylean are at an increased risk for exhibiting the downer pig syndrome (could also be referred to as leg weakness syndrome). Pig handling methods to reduce the incidence of downer pigs should be thoroughly evaluated prior to initiating use of Paylean.

WARNING: Not for use in humans. Individuals with cardiovascular disease should exercise special caution to avoid exposure. The active ingredient in Paylean, ractopamine hydrochloride, is a beta-adrenergic agonist.

WITHHOLDING PERIOD: MEAT - DO NOT USE less than 12 hours before slaughter for human consumption.

EXPORT TRADE ADVICE – TREATED STOCK: Exporters need to comply with the regulations/standards of importing countries with regard to the use of animal health products in livestock. Producers are advised to contact their export slaughter facility for information before giving pigs feed to which this product has been added.

SAFETY DIRECTIONS: Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. If product in eyes, wash it out immediately with water. Wash hands after use. When opening the container and preparing stock feed wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and a washable hat, elbow length PVC gloves and face shield or goggles. After each day's use, wash gloves and contaminated clothing

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 13 1126.

DISPOSAL OF BAGS: Shake and empty contents into medicated feed. Do not dispose of undiluted chemicals on site. Puncture or shred and bury empty bags in a local authority landfill. If not available bury the bag below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty bags and product should not be burnt.

Additional information is in the Material Safety Data Sheet.

Manufactured in U.K. for:
ELANCO ANIMAL HEALTH
 A division of Eli Lilly Australia Pty Limited A.B.N. 39 000 233 992
 112 WHARF ROAD, WEST RYDE, N.S.W. 2114
 TEL: Toll free 1800 226 324

APVMA Approval Number: 54172/10/1205

STORAGE: Store below 30°C (Room temperature) in a dry place.

Product Name	File Name	Version	Revision Date	Page No.
Paylean 20	Paylean 20 54172 1205 for Infopest.doc	Created on 3/14/2006 4:27:00 PM	Last printed 3/14/2006 4:31:00 PM	Page 2 of 3

For batch number and expiry date see reverse

USE NO HOOKS

Bar code 93 23829 00273 6

* Elanco[®], Paylean[®] and the diagonal colour bar are trademarks of Eli Lilly and Company

Product Name	File Name	Version	Revision Date	Page No.
Paylean 20	Paylean 20 54172 1205 for Infopest.doc	Created on 3/14/2006 4:27:00 PM	Last printed 3/14/2006 4:31:00 PM	Page 3 of 3