

**PRESCRIPTION ANIMAL REMEDY**  
KEEP OUT OF REACH OF CHILDREN  
FOR ANIMAL TREATMENT ONLY



AH0635  
**ELANCO**

**Apralan**  
Soluble Powder

Store below 30°C (Room Temperature)  
in a dry place.

NRA No. 47742/01

Expiry Date:

Batch Number:

YL9000DEALX YL9000DEALX Manufactured in U.K.

Indications:

**Pigs:** For the treatment of bacterial enteritis caused by susceptible strains of *Escherichia coli* and other organisms susceptible to apramycin

**Calves:** For the prophylaxis or treatment of bacterial enteritis in calves caused by *E. coli* and *Salmonella* organisms susceptible to apramycin.

**Broilers:** For use in the control and treatment of colisepticaemia (colibacillosis) in broilers caused by susceptible strains of *E. coli*.

Active Constituent: 50g APRAMYCIN  
(as apramycin sulfate)

Contents: 50g APRAMYCIN as apramycin sulfate

Minimum 84g NET

Elanco Animal Health  
A Division of Eli Lilly Australia Pty. Limited  
A.C.N. 000 233 992  
112 Wharf Road, West Ryde, N.S.W. 2114  
Tel: (02) 9325 4455 or 1800 226 324

**DIRECTIONS FOR USE:**

**Do not administer for a period longer than 14 days.**

**READ ATTACHED LEAFLET BEFORE USING THIS PRODUCT FOR DOSAGE AND DIRECTIONS FOR USE.**

**WITHHOLDING PERIOD: MEAT - DO NOT USE less than 14 days before slaughter for human consumption.**

**DISPOSAL:** Dispose of empty container by wrapping with paper and putting in garbage.

Run finger along perforated edge, pull out leaflet

PA9500DEALP

---

PRESCRIPTION ANIMAL REMEDY  
KEEP OUT OF REACH OF CHILDREN  
FOR ANIMAL TREATMENT ONLY

**Apralan**  
Soluble Powder

Active Constituent: 50g **APRAMYCIN**  
(as apramycin sulfate)

**Indications**

**Pigs:** For the treatment of bacterial enteritis caused by susceptible strains of *Escherichia coli* and other organisms susceptible to apramycin

**Calves:** For the prophylaxis or treatment of bacterial enteritis in calves caused by *E. coli* and *Salmonella* organisms susceptible to apramycin.

**Broilers:** For use in the control and treatment of colisepticaemia (colibacillosis) in broilers caused by susceptible strains of *E. coli*.

**DIRECTIONS FOR USE**

**Do not administer for a period longer than 14 days.**

NOTE: THE BOTTLE IS PROVIDED WITH A MEASURING CAP.

**CAUTION:** Drinking water and delivery system must be free of rust to avoid reduction in activity.

**1. PIGS**

**Dosage**

**Neonatal:** Pigs should receive 25mg apramycin activity per kg of body weight daily. Start treatment on the day of birth and continue once daily for two additional days.

**Weanling:** For treatment of colibacillosis, pigs should consume sufficient medicated drinking water to obtain 12.5mg of apramycin activity per kg of body weight daily. Continue treatment for seven days. Prepare fresh medication solution daily to assure activity.

## Leaflet (Cont'd)

### **Mixing Directions - Pigs**

**Neonatal:** To prepare solution for treatment of neonatal pigs, **fill the smaller measure** (approx. 1g active ingredient), **add to 25mL of water** and mix well. When completely dissolved, **administer 1mL** of the solution **to each pig**.

**Weanling:** Add APRALAN Soluble Powder to the only source of drinking water to provide desired apramycin activity per litre. See table below.

To mix effectively, add the appropriate quantity of APRALAN Soluble Powder to five times its own volume of water. Stir well and allow **solution to stand for a few minutes then stir again**. When completely dissolved, add to the full volume of water and stir well. Prepare fresh medicated water every 24 hours to assure activity.

Approx. total body Wt <sup>a</sup> to be treated (kg)	Approx. daily water consumption (litres)	Apralan Soluble Powder	Apramycin Activity
80	10	1 small measure	1g
400	50	1 large measure	5g
4000	500	Entire Bottle	50g

<sup>a</sup> Number of pigs X average weight.

## **2. CALVES**

### **Dosage**

**Treatment:** Calves should receive 20 to 40mg of apramycin activity per kg of body weight daily according to the severity of the disease.

For **prophylaxis**, calves should receive 10mg apramycin activity per kg body weight daily according to the severity of the disease.

Continue treatment for 5 days. Add APRALAN Soluble Powder to the drinking water, milk or replacer according to the table below.

Leaflet (Cont'd)

Directions for use (Cont'd)

### **Mixing Directions - Calves**

Add the appropriate quantity of APRALAN Soluble Powder to five times its own volume of water. **Stir well and allow solution to stand a few minutes then stir again.** When completely dissolved, add to the full volume of drinking water, milk or milk replacer.

Solutions should be prepared immediately before use to assure activity.

Total body weight <sup>a</sup> (kg)		APRALAN Soluble Powder	Apramycin Activity
100	Prophylaxis	1 small measure	1g
	Treatment	2-4 small measures	2-4g
500	Prophylaxis	1 large measure	5g
	Treatment	2-4 large measures	10-20g
2,500	Prophylaxis	5 large measures	25g
	Treatment	1-2 bottles	50-100g

<sup>a</sup> Number of calves X average weight.

### **3. BROILERS**

#### **Dosage**

250 to 500mg apramycin activity/litre of drinking water (one to two bottles per 200 litres) for 5 consecutive days. This is equivalent to an average of 40-80mg apramycin activity per kg bodyweight daily.

#### **Mixing Directions - Broilers**

Add the appropriate quantity of APRALAN Soluble Powder to five times its own volume of water. **Stir well and allow solution to stand for a few minutes then stir again.** When completely dissolved, add to the full volume of drinking water and stir well. Prepare fresh medicated drinking water every 24 hours to assure activity.

**WITHHOLDING PERIOD: DO NOT USE LESS THAN 14 DAYS BEFORE  
SLAUGHTER FOR HUMAN CONSUMPTION.**

Leaflet (Cont'd)

**Disposal:** Dispose of empty container by wrapping with paper and putting in garbage.

**Storage:** Store below 30°C (Room Temperature) in a dry place.

ELANCO ANIMAL HEALTH  
A DIVISION OF ELI LILLY AUSTRALIA PTY. LIMITED  
A.C.N. 000 233 992  
112 WHARF ROAD, WEST RYDE, N.S.W. 2114  
TEL: (02) 9325 4455 OR 1800 226 324

NRA Approval No. 47742/01

Manufactured in U.K.