

**PRESCRIPTION ANIMAL REMEDY**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**  
**FOR ANIMAL TREATMENT ONLY**

ELANCO AF0472



Premix

ACTIVE CONSTITUENT: 200g/kg TILMICOSIN (as the phosphate)

For the control of pneumonia in swine, caused by *Actinobacillus pleuropneumoniae*, *Mycoplasma hyopneumoniae*, *Pasteurella multocida*, and other organisms sensitive to tilmicosin.

**FOR USE IN SWINE FEEDS ONLY**  
**DO NOT FEED UNDILUTED**

**10 kg NET**

**DIRECTIONS FOR USE:**

**Do not allow horses or other equines access to feeds containing tilmicosin.**

**The safety of tilmicosin has not been established in boars intended for breeding purposes.**

**Do not use PULMOTIL 200 Premix in any feed (supplement, concentrate or complete) containing bentonite.**

**Bentonite in feeds may affect the efficacy of tilmicosin.**

**Any variation to the registered dose, frequency, duration, route, disease or target species may result in the need to extend the registered withholding period.**

Tilmicosin Concentration in Complete Feed (ppm*)	Kg of Pulmotil 200 Premix per 1000 kg of Complete Feed
200	1kg
400	2kg

(\* ppm = parts per million)

PRODUCT NAME:	FILE NAME:	REVISION NO. & DATE	LABEL CODE	
Pulmotil 200 Premix	46076_0501c.doc	1.1, 11 <sup>th</sup> May 2001	TBA	Page 1 of 2

**Mixing directions:**

IMPORTANT - PULMOTIL 200 PREMIX MUST BE THOROUGHLY MIXED IN FEED BEFORE USE.

Thoroughly mix PULMOTIL 200 Premix into the ration as outlined below. To ensure adequate mixing, it is recommended that a pre-blend of PULMOTIL 200 Premix be made before incorporation into the total amount of complete feed.

**Feeding directions:**

Feed continuously as the sole ration for a 14 day period, beginning approximately 7 days before an anticipated disease outbreak. If clinical signs of pneumonia have already been observed, clinically ill pigs may receive supplementary treatment by injection of an approved antibiotic.

**WITHHOLDING PERIOD: REMOVE ALL MEDICATED FEED FOURTEEN (14) DAYS BEFORE SLAUGHTER FOR HUMAN CONSUMPTION.**

**SAFETY DIRECTIONS: Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. Do not inhale dust. When mixing wear elbow-length PVC gloves and goggles and if dust is present half-face respirator with dust cartridge or canister. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles, respirator, and if rubber wash with detergent and warm water, and contaminated clothing.**

**PROTECTION OF THE ENVIRONMENT:** Sludge from waste treatment ponds must be disposed of in approved landfills for contaminated wastes. This product may accumulate in soils. DO NOT spread effluent containing tilmicosin on the same pastures or croplands in successive years. DO NOT allow wastes from pigs treated with tilmicosin to enter waterways. These wastes contain tilmicosin which is highly toxic to algae.

**DISPOSAL OF BAGS:** Shake bag empty into medicated feed. Puncture and bury empty bags in a local authority landfill. If not available bury the bag below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty bags and product should not be burnt. Do not dispose of undiluted premix on site.

NRA No: 46076/0501

Member  
(logo) Avcare

Manufactured in U.K. for:  
ELANCO ANIMAL HEALTH  
A DIVISION OF ELI LILLY AUSTRALIA PTY. LIMITED  
A.B.N. 39 000 233 992  
112 WHARF ROAD, WEST RYDE, N.S.W. 2114  
TEL: 1800 226 324

Store below 30°C (Room Temperature) in a dry place.  
For expiry date and batch number see reverse.

# USE NO HOOKS

PRODUCT NAME:	FILE NAME:	REVISION NO. & DATE	LABEL CODE	
Pulmotil 200 Premix	46076_0501c.doc	1.1, 11 <sup>th</sup> May 2001	TBA	Page 2 of 2