

Printed label, code no. WS1740ALX

PRESCRIPTION ANIMAL REMEDY
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY

AH0206

ELANCO

Tylan

200

Tylosin injection
200mg/mL



ACTIVE CONSTITUENT: 200mg/mL TYLOSIN

FOR THE TREATMENT OF TYLOSIN SUSCEPTIBLE
BACTERIAL AND MYCOPLASMAL INFECTIONS IN CATTLE
AND PIGS

100mL

NRA Approval No: 36796/1101

Manufactured in U.S.A. for
ELANCO ANIMAL HEALTH
A DIVISION OF ELI LILLY AUSTRALIA PTY LIMITED
A.C.N. 000 233 992
112 WHARF ROAD, WEST RYDE, N.S.W. 2114
Tel: (02) 9325 4455 or 1800 226 324

**READ THE ACCOMPANYING LEAFLET BEFORE USING THIS
PRODUCT.**

DIRECTIONS FOR USE:

Dosage

CATTLE: 5-10mg tylosin per kg bodyweight daily i/m or slow i/v
(equivalent to 0.5-1mL Tylan 200 Tylosin Injection per 20kg
bodyweight). Do not exceed 5 days treatment.

PIGS: 5-10mg tylosin per kg bodyweight daily i/m (equivalent to
0.5-1mL Tylan 200 Tylosin Injection per 20kg bodyweight).
Do not exceed 3 days treatment.

WITHHOLDING PERIODS:

**MEAT: Cattle - DO NOT USE less than 21 days before slaughter for
human consumption.**

**MEAT: Pigs - DO NOT USE less than 3 days before slaughter for
human consumption.**

**MILK: Cattle - Milk taken from cows within 72 hours (6 milkings)
following treatment MUST NOT BE USED for human consumption
or processing.**

Disposal of containers and needles: Dispose of empty container by
wrapping with paper and putting in garbage. Discarded needles should
immediately be placed in a designated and appropriately labelled
"sharps" container.

Store below 25°C (Air Conditioning).

Expiry Date

Batch Number

YK8995AL

Printed carton, code no. SI6850ALS

Front panel



PRESCRIPTION ANIMAL REMEDY
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY



ACTIVE CONSTITUENT: 200mg/mL TYLOSIN

FOR THE TREATMENT OF TYLOSIN SUSCEPTIBLE BACTERIAL AND
MYCOPLASMAL INFECTIONS IN CATTLE AND PIGS

100mL

Manufactured in U.S.A. for:
ELANCO ANIMAL HEALTH
A DIVISION OF ELI LILLY AUSTRALIA PTY LIMITED
A.C.N. 000 233 992
112 WHARF ROAD, WEST RYDE, N.S.W. 2114.
Tel: (02) 9325 4455 or 1800 226 324

Side panel 1

READ THE ACCOMPANYING LEAFLET BEFORE USING THIS PRODUCT.

DIRECTIONS FOR USE

Dosage

Cattle: 5-10mg tylosin per kg bodyweight daily i/m or slow i/v (equivalent to 0.5-1mL Tylan 200 Tylosin Injection per 20kg bodyweight). Do not exceed 5 days treatment.

Pigs: 5-10mg tylosin per kg bodyweight daily i/m (equivalent to 0.5-1mL Tylan 200 Tylosin Injection per 20kg bodyweight). Do not exceed 3 days treatment.

WITHHOLDING PERIODS:

MEAT: Cattle - DO NOT USE less than 21 days before slaughter for human consumption.

MEAT: Pigs - DO NOT USE less than 3 days before slaughter for human consumption.

MILK: Cattle - Milk taken from cows within 72 hours (6 milkings) following treatment MUST NOT BE USED for human consumption or processing.

Any variation by the prescribing veterinarian to the approved dose, frequency, duration, route, disease, or target species may require extending the approved withholding period.

Printed carton, code no. SI6850ALS

Side panel 2

Additional information is in the Material Safety Data Sheet.

Disposal of containers and needles: Dispose of empty container by wrapping with paper and putting in garbage. Discarded needles should immediately be placed in a designated and appropriately labelled “sharps” container.

NRA Approval No: 36796/1101

Store below 25°C (Air Conditioning). Use all product within 90 days of opening and discard any unused portion.

Expiry Date

Batch Number

Printed carton, code no. SI6850ALS

Top panel



100mL

PRESCRIPTION ANIMAL REMEDY
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY

AH0206
ELANCO
Tylan 200
Tylosin injection 200mg/mL

INDICATIONS:

For the treatment of tylosin susceptible bacterial and mycoplasmal infections in cattle and pigs.

DESCRIPTION:

TYLAN 200 Tylosin Injection is an antibiotic substance produced by a strain of *Streptomyces fradiae*, an actinomycete. Tylan 200 Tylosin Injection is a sterile solution of tylosin base in 50% propylene glycol. Each millilitre contains 200 milligrams of tylosin base activity.

PROPERTIES:

Tylosin is used exclusively in animals. The oral LD₅₀ in mice is greater than 5 grams per kilogram. In pigs, the LD₅₀ by the intramuscular route is approximately 1000 milligrams per kilogram; by oral route, greater than 5 grams per kilogram. Blood level studies have been conducted in cattle and pigs comparing TYLAN 200 Tylosin Injection to aqueous solutions of tylosin tartrate. A longer duration of activity was evidenced with TYLAN 200 Tylosin Injection.

Tylosin has an antibacterial spectrum that is essentially gram-positive, but it is also active against certain spirochaetes, large viruses, and certain gram negative organisms (not including coliforms). It has been found to be extremely active against the Mycoplasma species.

INDICATIONS:

Cattle: TYLAN 200 Tylosin Injection is effective in cattle for the treatment of pneumonia, metritis, foot rot, leptospirosis, acute mastitis, contagious calf pneumonia, pneumoenteritis of calves, secondary bacterial infections associated with virus diseases, and as an aid in combating infections associated with surgical procedures or injuries. In general, this drug is effective against those conditions associated with or caused by bacteria sensitive to tylosin.

Pigs: TYLAN 200 Tylosin Injection is indicated as an aid in treating the secondary bacterial invaders in virus diseases. Pig diseases reported responding to TYLAN 200 Tylosin Injection therapy are: erysipelas, pneumonia and dysentery in pigs.

DIRECTIONS FOR USE

Use all product within 90 days of opening and discard any unused portion.

Dosage and administration:

Cattle: Inject 5 to 10mg tylosin per kilogram of bodyweight daily intramuscularly or by slow intravenous injection (which is equivalent to 0.5mL to 1mL Tylan 200 Tylosin Injection per 20 kg bodyweight). Treatment should be continued 24 hours following remission of disease signs. Do not exceed 5 days of treatment.

Pigs: Inject 5 to 10mg tylosin per kilogram of bodyweight daily intramuscularly (which is equivalent to 0.5mL to 1mL Tylan 200 Tylosin Injection per 20 kg bodyweight). Treatment should be continued 24 hours following remission of disease signs. Do not exceed 3 days of treatment.

Do not mix TYLAN 200 Tylosin Injection with other parenteral solutions as this may cause precipitation of the active ingredient.

CAUTION: If there is no response to therapy in three days, diagnosis and treatment should be reassessed.

CONTRA-INDICATIONS AND SIDE EFFECTS:

There are no known contra-indications to TYLAN 200 Tylosin Injection at this time. . Occasionally pigs have been observed with a reaction to TYLAN 200 Tylosin Injection. This consists of an oedema of the rectal mucosa, mild anal profusion, diarrhoea, erythema and pruritis. Discontinuation of treatment has effected an uneventful recovery from these symptoms. As with the injection of any foreign substance, an injection site reaction may very occasionally occur in some pigs. If the reaction is of concern to the producer, allowing a period of at least three weeks to elapse before slaughter will minimise the need for excessive trim at the injection site.

WITHHOLDING PERIODS:

MEAT: Cattle - DO NOT USE less than 21 days before slaughter for human consumption.

MEAT: Pigs - DO NOT USE less than 3 days before slaughter for human consumption.

MILK: Cattle - Milk taken from cows within 72 hours (6 milkings) following treatment MUST NOT BE USED for human consumption or processing.

DISPOSAL OF CONTAINERS AND NEEDLES: Dispose of empty container by wrapping with paper and putting in garbage. Discarded needles should immediately be placed in a designated and appropriately labelled "sharps" container.

STORAGE: Store below 25°C (Air Conditioning)

Use all product within 90 days of opening and discard any unused portion.

PRESENTATION: Multidose vial of 100mL. Each mL contains 200mg of tylosin.

NRA Approval No: 36796/1101

Manufactured in U.S.A. for:

ELANCO ANIMAL HEALTH

A DIVISION OF ELI LILLY AUSTRALIA PTY LIMITED

A.C.N. 000 233 992

112 WHARF ROAD, WEST RYDE, N.S.W. 2114.

Tel: (02) 325 4455 or 1800 226 324