

PRESCRIPTION ANIMAL REMEDY
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

* AF0250

ELANCO



Premix

*



ACTIVE CONSTITUENT: 250g/kg TYLOSIN as the phosphate

For use as an aid in stimulating growth and improving feed efficiency in pigs and as an aid in controlling enteric diseases caused by organisms susceptible to tylosin in pigs.

For the treatment and prevention of ileitis in pigs.

For reduction in the incidence of liver abscess in cattle.

Label image proudly supplied by



NOTE: This product is not for resale in this form and directions for use are given as guidelines only. This pack is intended for use by feed formulators for incorporation in medicated premixes and, when utilised as recommended by the formulator, will supply registered levels of tylosin in the feed.

25kg NET

APVMA Approval No: 36806/25AUS/0104

Manufactured in Australia by:
ELANCO ANIMAL HEALTH
A DIVISION OF ELI LILLY AUSTRALIA PTY. LIMITED
A.B.N. 39 000 233 992
112 WHARF ROAD, WEST RYDE, N.S.W. 2114
TEL: (Toll Free): 1800 226 324

* Elanco®, Tylan® and the diagonal colour bar are trademarks of Eli Lilly and Company

YK6018AL

Tylan 250
Tylosin phosphate Premix

DIRECTIONS FOR USE:

Mixing directions: IMPORTANT - TYLAN 250 PREMIX MUST BE THOROUGHLY MIXED IN FEED BEFORE USE AT MAXIMUM USE RATE OF 0.4KG TYLAN 250 PREMIX PER TONNE OF FINAL FEED

	<u>Amount of Tylan 250 Premix mixed per tonne of feed</u>	<u>Tylosin Equivalent per tonne of feed</u>
<u>A. PIGS</u>		
As an aid in controlling enteric diseases caused by organisms susceptible to tylosin:		
a) For young piglets, birth to 20kg and for breeding, growing and introduced pigs	0.4kg	100g
As an aid in prevention of swine dysentery:		
a) For growing pigs, birth to 20kg	0.4kg	100g
b) For breeding and growing pigs, 20kg to market	0.16kg	40g
For the treatment of ileitis. Feed for 21 days:	0.4kg	100g
For the prevention of ileitis:	0.16kg	40g
For use as an aid in stimulating growth and improving feed efficiency in pigs:		
a) Up to 20kg liveweight	0.4kg	100g
b) 20kg to 50kg liveweight	0.08kg	20g
c) 50kg to market	0.04kg	10g

B. CATTLE

For reduction in the incidence of liver abscess:

In Complete Feeds: 11 ppm tylosin, i.e. 0.044kg Tylan 250 Premix per tonne of feed.

Supplementary Feeds: Dependent on total dry matter intake of animal and supplementary feeding rate. Ensure that amount of Tylan fed does not exceed 11ppm of total dry matter intake (supplement plus all other feeds consumed per day). As an example, for dairy cows, addition according to the following table provides 150mg tylosin/head/day.

<u>Supplementary Feeding Rate kg/head/day</u>	<u>Amount of Tylan 250 Premix per tonne of feed</u>	<u>Tylosin equivalent per tonne of feed</u>
1	0.6 kg	150.0 g
2	0.3 kg	75.0 g
4	0.15 kg	37.5 g
6	0.1 kg	25.0 g
8	0.075 kg	18.75g



Make changes to diet gradually. Increase tylosin in feed up to desired dose over 5-7 days.

Any variation by the prescribing veterinarian to the approved dose, frequency, duration, route, disease or target species may require extending the approved withholding period.

WITHHOLDING PERIODS - MEAT: NIL (at 100g or less tylosin per tonne of final feed). MILK: NIL

SAFETY DIRECTIONS: Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. Repeated exposure may cause allergic disorders. Do not inhale dust. When opening the container and mixing into premixes wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow length PVC gloves and disposable dust mask. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 13 1126.

DISPOSAL OF BAGS: Shake and empty contents into medicated feed. Do not dispose of undiluted chemicals on site. Puncture or shred and bury empty bags in a local authority landfill. If not available bury the bag below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty bags and product should not be burned.

Additional information is in the Material Safety Data Sheet.

Store below 30°C (Room Temperature) in a dry place.

Expiry date:

Batch number:

USE NO HOOKS

YK6019AL

PRESCRIPTION ANIMAL REMEDY

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

ELANCO * AF0250

Tylan[®] 250



Tylosin Phosphate
Premix

*

ACTIVE CONSTITUENT: 250g/kg TYLOSIN as the phosphate.

**For use as an aid in stimulating growth and improving feed efficiency in pigs and as an aid in controlling enteric diseases caused by organisms susceptible to tylosin in pigs.
For the treatment and prevention of ileitis in pigs.**

For reduction in the incidence of liver abscess in cattle.

NOTE: This product is not for resale in this form and directions for use are given as guidelines only. This pack is intended for use by feed formulators for incorporation in medicated premixes and, when utilised as recommended by the formulator, will supply registered levels of tylosin in the feed.

25kg NET

APVMA Approval No: 36806/25UK/0104

Manufactured in UK for:
ELANCO ANIMAL HEALTH
A DIVISION OF ELI LILLY AUSTRALIA PTY. LIMITED
A.B.N. 39 000 233 992
112 WHARF ROAD, WEST RYDE, N.S.W. 2114
TEL: (Toll Free): 1800 226 324



*Elanco®, Tylan® and the diagonal colour bar are trademarks of Eli Lilly and Company

BG7240DEALL (I-MAY-04)
BGG550

Store below 30°C (Room Temperature) in a dry place.

Batch number:

Expiry date:

USE NO HOOKS

ELANCO * AF0250

Tylan® 250

Tylosin Phosphate Premix



MEMBER



DIRECTIONS FOR USE:

Mixing directions: IMPORTANT - TYLAN 250 PREMIX MUST BE THOROUGHLY MIXED IN FEED BEFORE USE AT MAXIMUM USE RATE OF 0.4KG TYLAN 250 PREMIX PER TONNE OF FINAL FEED

A. PIGS

As an aid in controlling enteric diseases caused by organisms susceptible to tylosin:

	<u>Amount of Tylan 250 Premix mixed per tonne of feed</u>	<u>Tylosin Equivalent per tonne of feed</u>
a) For young piglets, birth to 20kg and for breeding, growing and introduced pigs	0.4 kg	100 g

As an aid in prevention of swine dysentery:

a) For growing pigs, birth to 20kg	0.4 kg	100 g
b) For breeding and growing pigs, 20kg to market	0.16 kg	40 g

For the treatment of ileitis. Feed for 21 days:	0.4 kg	100 g
--	--------	-------

For the prevention of ileitis:	0.16 kg	40 g
---	---------	------

For use as an aid in stimulating growth and improving feed efficiency in pigs:

a) Up to 20kg liveweight	0.4 kg	100 g
b) 20kg to 50kg liveweight	0.08 kg	20 g
c) 50kg to market	0.04 kg	10 g

B. CATTLE

For reduction in the incidence of liver abscess:

In Complete Feeds: 11 ppm tylosin, i.e. 0.044kg Tylan 250 Premix per tonne of feed.

Supplementary Feeds: Dependent on total dry matter intake of animal and supplementary feeding rate. Ensure that amount of Tylan fed does not exceed 11ppm of total dry matter intake (supplement plus all other feeds consumed per day). As an example, for dairy cows, addition according to the following table provides 150mg tylosin/head/day.

Supplementary Feeding Rate kg/head/day	Amount of Tylan 250 Premix per tonne of feed	Tylosin equivalent per tonne of feed
1	0.6 kg	150.0 g
2	0.3 kg	75.0 g
4	0.15 kg	37.5 g
6	0.1 kg	25.0 g
8	0.075 kg	18.75 g

Make changes to diet gradually. Increase tylosin in feed up to desired dose over 5-7 days.

Any variation by the prescribing veterinarian to the approved dose, frequency, duration, route, disease or target species may require extending the approved withholding period.

WITHHOLDING PERIODS - MEAT: NIL (at 100g or less tylosin per tonne of final feed).

MILK: NIL

SAFETY DIRECTIONS: Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. Repeated exposure may cause allergic disorders. Do not inhale dust. When opening the container and mixing into premixes wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow length PVC gloves and disposable dust mask. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre.

Phone 13 1126.

DISPOSAL OF BAGS: Shake and empty contents into medicated feed. Do not dispose of undiluted chemicals on site. Puncture or shred and bury empty bags in a local authority landfill. If not available bury the bag below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty bags and product should not be burned.

Additional information is in the Material Safety Data Sheet.